



M. Dy.

Flore de l'Auvergne
Champs et jardins

E 40 - 16

1899

Carex hirta. (Linné.)
Carcassonne, bords de l'Aude.



Carex montana. (Linné)
Gaunes.

Carex montana



Carex *muricata.* (Linné)
Caracasenne *prés.*

Carex

pseudo-arenaria.
Montolieu.
(Reichenbach)





Carex pullicaris. (Linné.)
Montdieu.



Carex setifolia. (Godron)
Marseillette.



Carex stellulata. (Goodenough)
Carcassonne, bords du canal.



Eleocharis ovata. (Robert Brown)
Vimoux.



Schæenus
Environs

nigricans. (Linné.)
de La Franqui.



Agileps triuncialis. (Linné.) (Cépée)



Agrostis elegans. (Thores)
 Malepère aux Jasses.

Agrostis

Limoux

interrupta. (Linné.)

Lieux salinés.





Agrostis rupestris. (allioni.)
Belcain.



Agrostis vulgaris. (Withering.)
L'espunassière, bois.



Aira cypraniana. (Gussone.)
Carcassonne.

A. cypraniana



Aira elegans. (Gaudin.)
Pradelles Cabardès.



Brachypodium pinnatum. (S. de Beauvois)
Carcassonne lieux incultes.



Bromus erectus. (Hudson)
Saint-Jean, près de la Chaussée.



Bromus
det.

Madritensis. (Linné.)
lieux stériles.



Arundo Donax. (Linné)
Carcassonne



A detailed botanical illustration of the grass *Avena sesquiltertia*. The drawing shows two slender, green stems rising from a common base. Each stem is topped with a dense, elongated spike of small, light-colored flowers. The leaves are narrow and linear, with some showing a slight curve. The base of the plant features a cluster of dry, brownish roots or seed heads. The entire illustration is rendered in a classic, scientific style with fine lines and subtle color washes.

Avena *sesquiltertia.* (Linné.)
Gorges de Saint-Georges.

E. CHRYSON, ANCHUTHE, GAINSON & MONTGOMERY



Catapodium loliaceum. (Link.)
Leucate.

Catapodium loliaceum
Leucate



Cynosurus echinatus. (Linné.)
Carcassonne Bois de Serres.



Deschampsia caespitosa. (Paul Brown.)
-Lespinassière.



Deschampsia flexuosa. (Grisebach?)
Saint-Jeak.





Festuca elatior,
var. pratensis. (Hudson.)
Simoux, pres.



Festuca interrupta. (Desfontaines.)
Linnæus, Cords de St. André.



Festuca ovina, var. alpina. (Grenier et Godron.)
Forêt de La Tajeole.



Festuca rubra, (Linné)
Carcassonne, près.

Festuca rubra



Festuca
Alarie *spadicosa.* (Linné.)



Festuca varia, var. Flavescens. (Lapeyrouie.)
Gorges de Saint-Georges.



Gastridium lepidigerum. (Gaudin.)
Maleprie aux Jasses.



Gaudinia fragilis. (P. de Beauvois)
Pois de la Malepière.




Glyceria convoluta. (Fries)
He de Saute.



Glyceria fluitans. (Robert Brown)
Carcassonne, Fresquid.



Glyceria fluitans,
var. plicata. (Fries)
Montolieu.



Kæleria albescens. (de Candolle.)
La Clape.



Kæleria villosa. (Persoon.)
Sainte-Lucie



Lepturus cylindricus. (Vrin.)
Steud.



Lolium
limoux

arvense. (Schrader.)
champs.



Mibora verna. (Salisot de Beauvois)
Champs à Bouichère



Milium multiflorum. (Cavanilles)
Ile de Sainte-Lucie.



Milium paradoxum. Linné.
Fontfroide.



Nardurus tenellus. (Reich.)
Sabbie du Vaucluse.



Nardus stricta. (Linné.)
Forêt de la Fajede.



Phalaris paradoxa. (Linné.)
Simoux lieux cultivés.



Phleum Alpinum. (Linné.)
Forêt de La Fajede.



Poa compressa. (Linné.)
Carcassonne, Terrains secs.



Stipa

pennata. (Linné.)
Coteaux d'Allet.



Crisetum flavescens. (P. de Beauv.)
Barcassonne, talus des fossés du canal.



FLORE
DE
L'AUDE

Jardins



Asperula hirta. (Ramond.)



Crocus minimus. (DeCandolle)



Deutzia scabra. (Thunberg.)



Eryngium Alpinum. (Linné.)



Escallonia rubra. (Persoon.) *rubra*



Fagus cuprea.



Fagus purpurea.



Galega officinalis. (Linné.)



Helianthus multiflorus. (Linné.)



Humulus Japonicus. (Siebold et Zuccarini)
Variété foliis variegatis



Hypericum pratense.



Jasminum odoratissimum. (Linné.)
Jasminum officinale



Juniperus Virginiana. (Linné.)



Ligustrum Japonicum. (Thunberg.)



Sinaria bijartita. (Willdenow.)



Malope trifida. (Cavanilles.)



Malus communis. (Lamarch.)



Matthiola Græca. (Sweet.)



Mesembrianthemum cordifolium. (Linné.)



Mina lobata. (Sallemant et Exarca)



Muscari betryoides. (Miller.)



Negundo fraxinifolium. (Walt.)
variété variegata-nivea.



Nerium Oleander. (Linné.)

A detailed botanical illustration of Ornithogalum nutans. The plant features a central, upright stem with several long, narrow, lanceolate leaves that are green with prominent parallel veins. At the top of the stem is a dense, elongated inflorescence (raceme) bearing numerous small, bell-shaped flowers. The flowers are light-colored, possibly white or pale pink, with darker markings on the petals. The entire illustration is set against a plain, light-colored background.

Ornithogalum nutans. (Linné.)



Papaver somniferum. (Linné.)



Pelargonium grandiflorum. (Hort.)



Petunia violacea. (Lindley.)



Petunia oculata. (Hort.)
Variété à fleurs oculées.



Platanus occidentalis. (Linné.)



Peonia fragrans. (Anders.)



Paeonia officinalis. (Wetz.)



Paeonia officinalis. (Wetz.)



Pontederia

cordata. (Linné.)



Prunus domestica. (Linné.)
Variété.



Prunus Lauro-Cerasus. (Linné.)



Pyrethrum

frutescens. (Willedenow)



Quercus torva. (Rose.)



Rosa alpina. (Linné.)
Variété *nuda.* (Grenier et Godron.)



Ruscus racemosus. (Linné.)



Scabiosa graminifolia. (Linné)



Statice Armeria (Smith.)



Styphnolobium
japonicum. (Schott)



Taxodium distichum. (Richard.)



Thunbergia alata. (Bojer.)



Viola alpestris. (Jordan.)



Zizyphus vulgaris. (Lamarck.)